

Positive Mass from Holographic Causality ^{*}

Don N. Page^{a,†}, Sumati Surya^{a,b,§} and Eric Woolgar^{b,a,¶}

^a Theoretical Physics Institute, University of Alberta
Edmonton, AB, Canada T6G 2G1

^b Dept. of Mathematical Sciences, University of Alberta
Edmonton, AB, Canada T6G 2G1

Abstract

For $n+1$ dimensional asymptotically AdS spacetimes which have holographic duals on their n dimensional conformal boundaries, we show that the imposition of causality on the boundary theory is sufficient to prove positivity of mass for the spacetime when $n \geq 3$, without the assumption of any local energy condition. We make crucial use of a generalization of the time-delay formula calculated in gr-qc/9404019, which relates the time delay of a bulk null curve with respect to a boundary null geodesic to the Ashtekar-Magnon mass of the spacetime. We also discuss holographic causality for the negative mass AdS soliton and its implications for the positive energy conjecture of Horowitz and Myers.

Positive energy theorems play a central role in gravitational theories, since they determine the classical stability of physical spacetimes within given asymptotic classes. Indeed, alternative versions of the proof of positivity exist, using surprisingly distinct techniques. In the asymptotically flat case, among the many approaches, one has the proof of positivity using minimal surfaces, due to Schoen and Yau [1], Witten's supersymmetry-inspired proof using spinorial techniques [2], Geroch's approach using the inverse mean curvature flow [3, 4] and Penrose, Sorkin and Woolgar's approach using the focusing of null geodesics near conformal infinity $\mathcal{I}$ [5]. Similarly, in the asymptotically AdS case one has the spinorial proof due to Abbott and Deser [6], and a proof based on null focusing techniques in [7].

In this paper, we present a new proof of positivity of energy for asymptotically AdS spacetimes based on the assumption of holography [8, 9, 10, 11]. While the AdS/CFT correspondence [12, 13, 14, 15] is a particular realization of holography, our result does not depend on the specifics of the holographic correspondence.

^{*}Alberta-Thy-08-02, hep-th/0204198

[†]Fellow, Canadian Institute for Advanced Research

[‡]email: don@phys.ualberta.ca

[§]e-mail: ssurya@phys.ualberta.math.ca

[¶]e-mail: ewoolgar@math.ualberta.ca

In holographic theories, causally related events in the $n + 1$ dimensional bulk theory have interpretations in the n dimensional holographic dual theory and vice versa. Questions of causality in holographic theories have been discussed in many papers [16, 17, 18, 19, 20, 21, 22, 23, 24]. For example, Kleban, McGreevy and Thomas [24] propose a relationship between causality in the holographic dual theory and matter energy conditions in the bulk.

We show that the requirement of causality in the holographic dual theory is in fact sufficient to prove that a certain weighted time-average of the mass of the spacetime is non-negative, given the assumption of a holographic correspondence, but without the assumption of any energy conditions. In the special case when the matter flux on the conformal boundary $\mathcal{I}$ vanishes, this means that the instantaneous mass itself is non-negative. Our proof is based on a generalization to arbitrary spacetime dimensions $n + 1 \geq 4$ of an expression for time delay in [7], which relates the time difference between a null geodesic on the conformal boundary $\mathcal{I}$ and a nearby bulk null curve to the Ashtekar-Magnon mass.

As in the AdS/CFT correspondence, we shall take a holographic correspondence to imply that certain fields in the (conformally completed) bulk theory, evaluated at a point on the conformal boundary, correspond to operators in the dual, or “boundary”, theory localized at the same point. We further assume *holographic causality*: if two points on the conformal boundary are causally related via the bulk causal structure, then there is a causal relation (e.g. non-commutation) between the two corresponding operators of the dual boundary theory. The boundary theory is thus defined to be causal only if these two operators are localized at points that are causally related in the conformal boundary causal structure. More crudely, causality implies that a signal cannot go through the bulk faster than it can go along the conformal boundary.

We begin by considering the pair: a point S on $\mathcal{I}$, dubbed the source, and an observer world-line R on $\mathcal{I}$, dubbed the receiver. Let Δt be the time taken by a fastest null geodesic γ to go from S to R along $\mathcal{I}$ and $\Delta t'$ the time taken by a bulk null curve γ' to go from S to R . If $\Delta t > \Delta t'$, this means that R will receive a signal from S via the bulk along γ' before it receives the fastest boundary signal γ , thus violating holographic causality. Holographic causality therefore requires that the time delay $\Delta T = \Delta t' - \Delta t$ between the fastest boundary null geodesic and any bulk null curve must always be non-negative [24].

When the spacetime is pure AdS, the time delay between the fastest boundary signal and any bulk signal is non-negative. It vanishes when the bulk signal is along a bulk null geodesic, but this can only join antipodal points on the boundary. Since the strictest version of AdS/CFT holography holds for pure AdS bulk spacetimes (in 3, 5 and 7 spacetime dimensions), holographic causality is satisfied. However, one must also be able to interpret, in the boundary theory, bulk perturbations about the global AdS background [25, 16]. Such perturbations would affect this exact matching and hence could lead to a violation of holographic causality. Intuitively one would expect that if global AdS is perturbed by the addition of positive mass, then causal geodesics in the bulk will focus and hence be non-maximal, thus maintaining holographic causality. In fact what we will find is the converse of this intuition — namely, the fact that bulk causal curves are non-maximal implies positivity of mass.




Figure 1: In order to preserve holographic causality, the time delay $\Delta T = \Delta t' - \Delta t$ must be non-negative.

To make our treatment self contained, we start by reminding the reader of the conformal definition of an asymptotically AdS spacetime in $n + 1$ spacetime dimensions for $n + 1 \geq 4$ [26, 27]; we use the definition in [27] rather than the weaker one in [26].

Definition: An $n + 1$ dimensional spacetime (M, g_{ab}) is said to be asymptotically AdS if there exists a manifold $\hat{M}$ with conformal boundary $\mathcal{I}$, with a (sufficiently) smooth metric $\hat{g}$ and a diffeomorphism from M to $\hat{M} - \mathcal{I}$ such that

- (i) There exists a function Ω on $\hat{M}$ such that $\hat{g}_{ab} = \Omega^2 g_{ab}$ on M .
- (ii) $\mathcal{I} = \partial \hat{M}$ has topology $S^{n-1} \times \mathbb{R}$, $\Omega = 0$ on $\mathcal{I}$ and $\nabla_a \Omega$ is nowhere vanishing on $\mathcal{I}$.
- (iii) g_{ab} satisfies $R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = 8\pi T_{ab}$ where $\Lambda < 0$ and $\Omega^{1-n} T_{ab}$ admits a smooth limit to $\mathcal{I}$,
- (iv) The Weyl tensor of g_{ab} is such that $\Omega^{3-n} \hat{C}_{abcd}$ is smooth on $\hat{M}$ and vanishes at $\mathcal{I}$.
- (v) For $n = 3$ only (where it does not follow from the other conditions) we additionally require that the conformal isometry group of $\mathcal{I}$ is the AdS group.

We confine our attention to the conformal metric $\hat{g}_{ab} = \Omega^2 g_{ab}$ in a small neighbourhood $\mathcal{N} = \mathbb{R} \times S^{n-1} \times [0, \epsilon] = \mathbb{R} \times \mathcal{U}$ of $\mathcal{I}$, where $\Omega \in [0, \epsilon]$. It is convenient to express $\hat{g}_{ab}$ to order $\mathcal{O}(\Omega^{n+1})$ in a coordinate frame $x = (\Omega, \vec{x})$ adapted to the level surfaces of Ω , where $\vec{x} = (t, \vec{\omega}_{n-1})$ are the coordinates in constant Ω hypersurfaces and $\vec{\omega}_{n-1}$ are the coordinates on the S^{n-1} sphere. The conformal metric in $\mathcal{N}$ may be expressed as [28, 29, 30]

$$d\hat{s}^2 = d\hat{s}_o^2 + \Omega^n f_{ab}(\vec{x}) dx^a dx^b + \mathcal{O}(\Omega^{n+1}), \quad (1)$$

where

$$d\hat{s}_o^2 = \hat{g}_{ab}^\circ dx^a dx^b = \Omega^2 ds_{AdS}^2 = \frac{1}{1 - \Omega^2} d\Omega^2 - dt^2 + (1 - \Omega^2) d\omega_{n-1}^2, \quad (2)$$

with ds_{AdS}^2 the AdS metric, $d\omega_{n-1}^2$ the standard “round” metric on an $n - 1$ sphere and $f_{ab}(\vec{x})$ a degenerate symmetric tensor with $f_{a\Omega} = 0$. For simplicity we use units in which $\Lambda = -\frac{n(n-1)}{2}$.

As we now show, the trace-free part of f_{ab} is in fact proportional to the electric part E_{ab} of the Weyl tensor $\hat{C}_{abcd}$,

$$E_{ab}(\vec{x}) = \lim_{\rightarrow \mathcal{I}} \frac{1}{\Omega^{n-2}} \hat{C}_{acbd} \hat{\nabla}^c \Omega \hat{\nabla}^d \Omega. \quad (3)$$

Using the form of the metric (1), $\hat{C}_{abcd}$ can be simplified to

$$\begin{aligned} \hat{C}_{abcd} = & -2\hat{g}_{de}^{\circ} \hat{\nabla}_{[a}^{\circ} \mathcal{C}_{b]e}^e - \frac{4}{n-1} (\hat{\nabla}_{[d}^{\circ} \hat{g}_{c][b}^{\circ} \mathcal{C}_{a]e}^e + \hat{\nabla}_e^{\circ} \hat{g}_{[a][c}^{\circ} \mathcal{C}_{d]b}^e) \\ & - \frac{4}{n(n-1)} \hat{g}_{a[c}^{\circ} \hat{g}_{d]b}^{\circ} \hat{g}^{\circ ef} \hat{\nabla}_{[e}^{\circ} \mathcal{C}_{m]f}^m + \mathcal{O}(\Omega^{n-1}), \end{aligned} \quad (4)$$

where $\hat{\nabla}_a^{\circ}$ is the connection compatible with the conformal AdS metric $\hat{g}_{ab}^{\circ}$, $\hat{\nabla}_a$ is the connection compatible with (1), the tensor $\mathcal{C}_{ab}^c$ is given by $\mathcal{C}_{ab}^c v_c = \hat{\nabla}_a^{\circ} v_b - \hat{\nabla}_b v_a$, and we have used the fact that AdS has vanishing Weyl curvature, $\hat{C}_{abcd}^{\circ} = 0$. Using (1), (2) and (3), and denoting the induced metric on $\mathcal{I}$ by $\hat{q}_{ab}^{\circ}$, we get

$$E_{ab} = -\frac{n(n-2)}{2} f_{ab} + \frac{(n-2)}{2} \hat{q}_{ab}^{\circ} (\hat{q}^{\circ cd} f_{cd}). \quad (5)$$

Reinserting this into (1), we obtain

$$d\hat{s}^2 = (1 + \Omega^n H(\vec{x})) d\hat{s}_{\circ}^2 - \frac{2}{n(n-2)} \Omega^n E_{ab}(\vec{x}) dx^a dx^b - \Omega^n H(\vec{x}) d\Omega^2 + \mathcal{O}(\Omega^{n+1}), \quad (6)$$

where $n H(\vec{x}) \equiv \hat{q}^{\circ kl} f_{kl}$ and where we have used the fact that $E_{a\Omega} = 0$. The definition of asymptotically AdS spacetimes above, from [27], implies $H(\vec{x}) = 0$, so our proof actually applies to the broader class of metrics given by (1).

We now choose S and R so that they project to the antipodes of the spatial boundary sphere $S^{n-1} \subset \mathcal{I}$. Next, we use the arc length $\sigma \in [0, \pi]$ to parameterise the null curves, where $d\sigma^2 = (1 - \Omega^2)^{-1} d\Omega^2 + (1 - \Omega^2) d\omega_{n-1}^2$ is the “round” metric on S^n with the equator at $\Omega = 0$. Let $\gamma : [0, 1] \rightarrow \mathcal{I}$ be one of the boundary null geodesics between S and R and let $\gamma' : [0, 1] \rightarrow \mathcal{N}$ be a bulk null curve which is a particular variation of γ with $\gamma'(0) = \gamma(0)$. Since $d\sigma^2$ is the round metric on S^n , γ projects to the equator at $\Omega = 0$, and we may now choose γ' to project to a nearby geodesic of $d\sigma^2$ in $\mathcal{U}$. This is always possible on the round S^n , since geodesics with the same starting point will focus at the antipodal point. Thus, for the varied curve γ' , $\Omega(\sigma) = \Omega_{\mathbf{m}} \sin(\sigma)$, where $\Omega_{\mathbf{m}}$ is the farthest that γ' ventures into the bulk. For a sufficiently small $\Omega_{\mathbf{m}} > 0$ we then obtain the following expression for the time delay in γ' with respect to γ :

$$\Delta T = -\frac{\Omega_{\mathbf{m}}^n}{n(n-2)} \int_0^{\pi} \sin^n(\sigma) E_{ab}(\vec{x}) \gamma^a \gamma^b d\sigma + \mathcal{O}(\Omega_{\mathbf{m}}^{n+1}), \quad (7)$$

where $\vec{x}$ and $\gamma^a = \frac{dx^a}{d\sigma}$ are functions of σ along the boundary null geodesic γ . The requirement of holographic causality implies a non-negative time delay, so that the right hand side of (7) must be non-negative. We now show that the integrand is related to the Ashtekar-Magnon mass.

The Ashtekar-Magnon(AM) mass is defined as [26, 27]

$$\mu[C] := -\frac{1}{8\pi(n-2)} \int_C E_{ab} \chi^a \chi^b dS, \quad (8)$$

where C is a space-like cut of $\mathcal{I}$, dS the associated volume element and χ^a a conformal timelike Killing vector on $\mathcal{I}$. In the coordinate frame $\vec{x} = (t, \vec{\omega}_{n-1})$ we use above, a natural choice is $\chi^a = (1, 0, \dots, 0)$ up to normalization, with C the round S^{n-1} sphere with volume element $d\Theta$, so that

$$\mu(t) = -\frac{1}{8\pi(n-2)} \int E_{00} d\Theta. \quad (9)$$

Now, up to a scale factor, on $\mathcal{I}$ the “spray” of future directed null geodesics at a point is parameterised by an $n-2$ dimensional sphere. Thus, for example, in $n+1=4$ dimensions there is a circle worth of null directions on $\mathcal{I}$ at every point, while in $n+1=5$ dimensions there is a 2-sphere worth of null directions. Taking $d\Xi$ as the volume element on the round S^{n-2} , we then take the average

$$-\int E_{ab} \gamma^a \gamma^b d\Xi = -\frac{n}{n-1} \text{vol}(S^{n-2}) E_{00}. \quad (10)$$

Here we have used (a) the fact that the integral is performed over the future null directions *at a point*, so that E_{ab} is in fact independent of these directions and (b) the fact that E_{ab} is trace-free. Averaging the time-delay (7) over all possible null directions as in (10) and over all points on the boundary S^{n-1} , we get

$$\int \Delta T d\Xi d\Theta = \frac{8\pi \text{vol}(S^{n-2})}{(n-1)} \Omega_{\mathbf{m}}^n \int_{t_0}^{t_0+\pi} \sin^n(t-t_0) \mu(t) dt + \mathcal{O}(\Omega_{\mathbf{m}}^{n+1}). \quad (11)$$

Since the time delay $\Delta T \geq 0$, this implies positivity of the weighted time-averaged mass,

$$\langle \mu \rangle_{t_0} = \int_{t_0}^{t_0+\pi} \sin^n(t-t_0) \mu(t) dt \geq 0. \quad (12)$$

If the matter flux $\int_V T_{ab} \chi^a \chi^b dV$ vanishes on $\mathcal{I}$ for the patch $V \subset \mathcal{I}$ bounded by spacelike cuts of $\mathcal{I}$ at t_0 and $t_0 + \pi$, then $\mu(t)$ is constant in V [26, 27], so that the instantaneous mass $\mu(t) \geq 0$ in V . We have thus proved the following:

Theorem: Assume that an $n+1$ dimensional asymptotically AdS spacetime (M, g) has a holographic dual on its conformal boundary $\mathcal{I}$. Then the requirement of holographic causality implies that the weighted time-averaged Ashtekar-Magnon mass $\langle \mu \rangle_{t_0}$ for the spacetime is non-negative for all $t_0 \in \mathbb{R}$. In particular, when the matter flux at $\mathcal{I}$ is zero for a time period of at least π , the instantaneous Ashtekar-Magnon mass is non-negative over that same time period.

Remark (i): In [7] the weight in $\langle \mu \rangle_{t_0}$ was assumed to be uniform, but we have shown here that it is indeed non-uniform. If this weight had been a delta function instead of $\sin^n(t-t_0)$, one could conclude that the instantaneous mass is non-negative in general. However, $\sin^n(t-t_0)$ only approaches a delta function as $n \rightarrow \infty$, so for finite n the weight has non-vanishing width. While this appears to be a drawback of the focusing method

employed here, one may be able to extract some useful information from it. In particular, (a) $\mu(t)$ cannot be negative for a time scale of $\sim \pi$, and (b) since $\langle \mu \rangle_{t_0} \geq 0$ for all t_0 , if $\mu(t) < 0$ on time-scales $\tau < \pi$, then it must be compensated for by a positive contribution from $\mu(t)$ for a time $\pi - \tau$ with “interest”. What does this say about classical stability? One might suppose that if the mass of the spacetime becomes negative at any time that this would lead to an increasingly negative mass. However, the fact that $\langle \mu \rangle_{t_0} \geq 0$ for all t_0 clearly prevents such an instability.

Remark (ii): The elements of our proof do not suffice to prove the converse of the theorem, namely that positivity of mass is sufficient to ensure holographic causality. Positivity of mass does not imply that the function $-E_{ab}\gamma^a\gamma^b$ is non-negative everywhere, and by choosing γ and γ' suitably, the time delay could be made negative. Instead of assuming positivity of mass however, if we were to assume null genericity, null geodesic completeness and a null energy condition, then a focusing theorem [31, 32, 33] can be used to show that all bulk geodesics have non-negative time-delay with respect to boundary null geodesics¹. This theorem states that in such spacetimes any complete causal geodesic must contain conjugate points. The existence of conjugate points means that the null geodesic focuses, so that the fastest null geodesics always lie on the boundary.

Remark (iii): In [24], it was claimed that any asymptotically AdS spacetime must satisfy an integrated weak energy condition if it is to have a causal holographic boundary dual. While the focusing theorem *implies* positive time delay, it is not clear how the converse conclusion of [24] may be reached.

Remark (iv): Finally, we briefly comment on the relation to the counterterm mass definition of [34], where the mass μ_{ct} is calculated by adding certain local counterterms to the action. In 4-dimensions it was shown in [27] that this mass definition is identical to the AM mass, so that holographic causality implies the positivity of μ_{ct} by our theorem. In 5 dimensions, however, these masses explicitly do not match, and their difference $\mu_{ct} - \mu$ can be interpreted as a non-zero Casimir energy. Using our method, however, holographic causality cannot be used to predict the sign of this Casimir energy term.

Now, the time delay formula (7) is valid only for asymptotically AdS spacetimes with $\mathcal{I}$ of topology $S^{n-1} \times \mathbb{R}$. However, asymptotically locally AdS spacetimes with non-spherical boundary are also candidates for a holographic theory [35, 36, 37, 38], and several examples of spacetimes with such boundaries are known [39]. Positive energy theorems have not been proved for such spacetimes — indeed, the AdS soliton [36], which has a Ricci flat $\mathcal{I}$, has negative AM mass². Motivated by the AdS/CFT correspondence, Horowitz and Myers [36] proposed a new positive energy conjecture which states that the AdS soliton is the unique ground state in its asymptotic class. It is therefore of interest to ask how the considerations in this current work extend to such spacetimes.

We now show by explicit calculation that despite having negative mass, the AdS soliton has a strictly positive time delay when $n \geq 3$, so that the boundary theory is causal in

¹The focusing theorem was used to complete the proof of positivity of energy in conjunction with the time delay formula in [7].

²The soliton is protected from a potential instability due to a unique scaling *isometry* between solitons of different mass parameters [40].

this restricted sense. The $n + 1$ dimensional AdS soliton has metric

$$ds^2 = -r^2 dt^2 + \frac{1}{V(r)} dr^2 + V(r) d\phi^2 + r^2 \sum_{i=1}^{n-2} (d\theta^i)^2. \quad (13)$$

Here $V(r) = \frac{r^2}{\ell^2} (1 - \frac{r_0^n}{r^n})$, with $\ell^2 = -\frac{n(n-1)}{2\Lambda}$, r_0 a constant and $n > 2$. The solution is non-singular provided ϕ is periodic with period $\beta_0 = \frac{4\pi\ell^2}{nr_0}$. Putting $x \equiv 1/r$, (13) can be re-expressed as

$$ds^2 = \frac{1}{x^2} (-dt^2 + ds_n^2), \quad (14)$$

where

$$ds_n^2 = \frac{\ell^2}{(1 - r_0^n x^n)} dx^2 + \frac{(1 - r_0^n x^n)}{\ell^2} d\phi^2 + \sum_i^{n-2} d\theta_i^2, \quad (15)$$

is the spatial part of the optical metric $d\hat{s}^2 = x^2 ds^2$. Thus, the time taken by a null signal is $\Delta t = \int ds_n$.

Now, consider a null geodesic from a source S on $\mathcal{I}$, which has angular momenta $(\vec{L}, J)$, where $\vec{L} \equiv (\dot{\theta}_1, \dots, \dot{\theta}_{n-2})$, $J \equiv (1 - r_0^n x^n) \dot{\phi}$ and $\vec{L} = \dot{J} = 0$ along any geodesic where $\dot{A} \equiv dA/ds_n$. The time taken for this geodesic to get back to $\mathcal{I}$ via the bulk (and *not* along the boundary) is then

$$\Delta t' = \sqrt{|\Delta \vec{\Theta}|^2 + \left(\frac{4\ell^2}{r_0^2} \right) (1 - k^2)^{\frac{2-n}{2n}} I(n)^2}, \quad (16)$$

where $|\Delta \vec{\Theta}|$ is the optical distance traversed in the $\vec{\Theta} \equiv (\theta_1, \dots, \theta_{n-2})$ directions, $k^2 = \ell^{-2} J^2 (1 - L^2)^{-1}$ and

$$I(n) = \int_0^1 \frac{1}{\sqrt{1 - X^n}} dX = \sqrt{\pi} \frac{\Gamma(\frac{n+1}{n})}{\Gamma(\frac{n+2}{2n})}. \quad (17)$$

If the bulk null geodesic is received by some observer R on $\mathcal{I}$, the shortest time taken by a boundary null geodesic to get from S to R satisfies the inequality,

$$\Delta t \leq \sqrt{|\Delta \vec{\Theta}|^2 + \left(\frac{4\ell^2}{r_0^2} \right) \frac{\pi^2}{n^2}}. \quad (18)$$

If the fastest curve, necessarily a null geodesic, were to lie in the bulk, then $\Delta t' < \Delta t$ for that geodesic. But we see that this cannot be the case. Comparing the times Δt and $\Delta t'$ we see that the time delay $\Delta T = \Delta t' - \Delta t$ is positive whenever

$$\frac{n}{\pi} I(n) (1 - k^2)^{\frac{2-n}{4n}} \geq 1. \quad (19)$$

Now, $(1 - k^2)^{\frac{2-n}{4n}} \geq 1$ for $n \geq 2$, the equality being satisfied only when $n = 2$ and/or $J = 0$. Moreover, one can easily show that $\frac{n}{\pi} I(n) \geq 1$ for $n \geq 2$ — indeed, it grows approximately linearly with n . Thus, the time-delay is zero only for $n = 2$. This is not surprising, since in 3 spacetime dimensions, the AdS soliton is in fact global AdS. It is

also satisfying to note that the existence of the time delay is in fact independent of the apparent mass “parameter” r_0 . This is consistent with the fact that solitons of different r_0 are in fact isometric to each other via a scaling isometry [40].

Thus, the AdS soliton, despite having negative mass, satisfies holographic causality. However, in light of the Horowitz-Myers conjecture, it is interesting that there is no bulk null geodesic which realizes the bound set by causality or alternatively, by the focusing theorem. What this means for the conjecture is a question under current investigation [41].

Acknowledgments: This work was partially supported by grants from the Natural Sciences and Engineering Research Council (Canada). SS was supported by a post-doctoral fellowship from the Pacific Institute for the Mathematical Sciences.

References

- [1] R. Schoen and S. T. Yau, Commun. Math. Phys. **65**, 45 (1979).
- [2] E. Witten, Commun. Math. Phys. **80**, 381 (1981).
- [3] R. Geroch, Ann. NY Acad. Sci. **224**, 108 (1973).
- [4] G. Huisken and T. Ilmanen, Int. Math. Res. Notices **20**, 1045 (1997).
- [5] R. Penrose, R. D. Sorkin and E. Woolgar, gr-qc/9301015.
- [6] L. F. Abbott and S. Deser, Nucl. Phys. B **195**, 76 (1982).
- [7] E. Woolgar, Class. Quant. Grav. **11**, (1994) 1881 [gr-qc/9404019].
- [8] L. Susskind, J. Math. Phys. **36**, 6377 (1995) [hep-th/9409089].
- [9] G. ’t Hooft, in *Salamfest*, eds. A. Ali, J. Ellis and S. Randjbar-Daemi, Singapore: World Scientific, 1994 [gr-qc/9310026].
- [10] D. Bigatti and L. Susskind, in *Strings, branes and gravity: Proceedings, Theoretical Advanced Study Institute, Boulder, USA, May 31-June 25, 1999*, eds. J. Harvey, S. Kachru, and E. Silverstein, River Edge: World Scientific, 2001 [hep-th/0002044].
- [11] R. Bousso, Rev. Mod. Phys. (in press) [hep-th/0203101].
- [12] J. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [hep-th/9711200].
- [13] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Phys. Lett. B **428**, 105 (1998) [hep-th/9802109].
- [14] E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [hep-th/9802150].
- [15] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, Phys. Rept. **323**, 183 (2000) [hep-th/9905111].
- [16] G. T. Horowitz and N. Itzhaki, JHEP **9902**, 010 (1999) [hep-th/9901012].
- [17] D. Kabat and G. Lifschytz, JHEP **9905**, 005 (1999) [hep-th/9902073].

- [18] D. Bak and S. J. Rey, Nucl. Phys. B **572**, 151 (2000) [hep-th/9902101].
- [19] S. B. Giddings, Phys. Rev. Lett. **83**, 2707 (1999) [hep-th/9903048].
- [20] J. Polchinski, L. Susskind and N. Toumbas, Phys. Rev. D **60**, 084006 (1999) [hep-th/9903228].
- [21] L. Susskind and N. Toumbas, Phys. Rev. D **61**, 044001 (2000) [hep-th/9909013].
- [22] J. Louko, D. Marolf and S. F. Ross, Phys. Rev. D **62**, 044041 (2000) [hep-th/0002111].
- [23] S. B. Giddings and M. Lippert, Phys. Rev. D **65**, 024006 (2002) [hep-th/0103231].
- [24] M. Kleban, J. McGreevy and S. Thomas, hep-th/0112229.
- [25] V. Balasubramanian, P. Kraus, A. E. Lawrence and S. P. Trivedi, Phys. Rev. D **59**, 104021 (1999) [hep-th/9808017].
- [26] A. Ashtekar and A. Magnon, Classical Quantum Gravity **1**, L39 (1984).
- [27] A. Ashtekar and S. Das, Class. Quant. Grav. **17**, L17 (2000) [hep-th/9911230].
- [28] M. Henneaux and C. Teitelboim, Commun. Math. Phys. **98**, 391 (1985).
- [29] C. Fefferman and C. R. Graham, Astérisque, hors série, p. 95 (1985); C. R. Graham, *Proc. 19th Winter School in Geometry and Physics*, Srni, Czech Rep., Jan. 1999.
- [30] M. Henningson and K. Skenderis, J. High Energy Phys. **9807**, 023 (1998) [hep-th/9806087].
- [31] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Spacetime*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 1973.
- [32] F. J. Tipler, J. Diff. Eq. **30** 165 (1978); Phys. Rev. D. **17** 2521 (1978).
- [33] A. Borde, Class. Quant. Grav. **4**, 343 (1987).
- [34] V. Balasubramanian and P. Kraus, Phys. Rev. Lett. **83**, 3605 (1999) [hep-th/9903190].
- [35] D. Birmingham, Class. Quant. Grav. **16**, 1197 (1999) [hep-th/9808032].
- [36] G. T. Horowitz and R. C. Myers, Phys. Rev. D **59**, 026005 (1999) [hep-th/9808079].
- [37] R. Emparan, C. V. Johnson and R. C. Myers, Phys. Rev. D **60**, 104001 (1999) [hep-th/9903238].
- [38] S. Surya, K. Schleich and D. M. Witt, Phys. Rev. Lett. **86**, 5231 (2001) [hep-th/0101134].
- [39] For a review see R.B. Mann, in *Internal Structure of Black Holes and Spacetime Singularities*, eds. L. Burko and A. Ori, Ann. Israeli Phys. Soc. **13**, 311 (1998) [gr-qc/9709039].

- [40] G. J. Galloway, S. Surya and E. Woolgar, Phys. Rev. Lett. **88**, 101102 (2002) [hep-th/0108170].
- [41] S. Surya and E. Woolgar (work in progress).